Arterial Thromboembolic Events Preceding the Diagnosis of Cancer in Older Persons

Babak B. Navi,¹⁻³ Anne S. Reiner,¹ Hooman Kamel,^{1,2} Costantino Iadecola,^{1,2} Peter M. Okin,⁵ Scott T. Tagawa,⁶ Katherine S. Panageas,⁴ and Lisa M. DeAngelis¹⁻³

¹Department of Neurology, Weill Cornell Medicine, New York, NY
 ²Feil Family Brain and Mind Research Institute, Weill Cornell Medicine, New York, NY
 ³Department of Neurology, Memorial Sloan Kettering Cancer Center, New York, NY
 ⁴Department of Epidemiology and Biostatistics, Memorial Sloan Kettering Cancer Center, New York, NY
 ⁵Department of Medicine, Division of Cardiology, Weill Cornell Medicine, New York, NY
 ⁶Department of Medicine, Division of Hematology and Medical Oncology, Weill Cornell Medicine, New York, NY

Correspondence/reprint requests to:

Babak B. Navi, MD, MS 525 East 68th Street, Room F610 New York, NY 10065 Phone: 212-746-0225 Fax: 646-962-0126 Email: ban9003@med.cornell.edu

Running head: Arterial Thromboembolism Before Cancer Diagnosis

Text word count: 3,709 Abstract word count: 250 References: 30 Tables: 4 Figures: 1 Scientific section designation: THROMBOSIS AND HEMOSTASIS

Key points

- Among 748,662 Medicare beneficiaries, the risk of arterial thromboembolic events was increased 69% in the year before cancer diagnosis.
- The increased risk of arterial thromboembolic events began 5 months before cancer was officially diagnosed and peaked in the month prior.

Abstract

Cancer patients face an increased risk of arterial thromboembolism; however, it's uncertain when this excess risk begins. This study evaluated the risk of arterial thromboembolism before cancer diagnosis. Using the population-based Surveillance Epidemiology and End Results-Medicare linked dataset, we identified 374,331 patients \geq 67 years of age with a new primary diagnosis of breast, lung, prostate, colorectal, bladder, uterine, pancreatic, gastric cancer, or non-Hodgkin lymphoma from 2005-2013. Cancer patients were individually matched by demographics and comorbidities to Medicare beneficiaries without cancer, who served as controls. Validated diagnosis codes were used to identify arterial thromboembolic events, defined as a composite of myocardial infarction or ischemic stroke. The Mantel-Haenszel estimator was used to compare risks of arterial thromboembolic events between cancer and non-cancer groups during 30-day periods in the 360 days before date of cancer diagnosis. From 360 to 151 days before cancer diagnosis, the 30-day interval risks of arterial thromboembolic events were similar between cancer patients and matched controls. From 150 to 1 day before cancer diagnosis, the interval 30day risks of arterial thromboembolic events were higher in cancer patients versus matched controls, progressively increasing as the cancer diagnosis date approached, and peaking during the 30-days immediately before cancer diagnosis, when 2,313 (0.62%) cancer patients were diagnosed with an arterial thromboembolic event versus 413 (0.11%) controls (odds ratio, 5.63; 95% confidence interval, 5.07-6.25). In conclusion, the risk of arterial thromboembolic events begins to increase 150 days before the date of cancer diagnosis in older persons and peaks in the 30 days before.

Introduction

An American's lifetime incidence of cancer is estimated to be 39%.¹ In the 6 months after diagnosis, cancer patients are twice as likely to develop myocardial infarction and ischemic stroke as compared to matched controls without cancer.² Some of this excess risk may be due to external factors associated with cancer, such as the effects of stress and cancer treatments, frequent interruption of antithrombotics, and increased surveillance.³ However, cancer itself, through effects on hemostatic and fibrinolytic pathways, may be the biggest driver of thromboembolism risk, as evidenced by the wide variability in risk by cancer type and stage.^{2,4-6} Cancer's prothrombotic effects include hematogenous release of cancer-derived microparticles which trigger the coagulation cascade, production of procoagulant factors such as factor X, release of mucins that activate platelets and endothelial cells through binding of P-selectin, and stimulation of neutrophils to release decondensed chromatin which form prothrombotic neutrophil extracellular traps.^{3,7,8} These effects, while linked to overall tumor burden, may precede cancer diagnosis, and thus it is speculated that some cancers can disrupt the body's hemostatic system and produce thrombosis months or even years before detection.⁹ For example, up to 10% of venous thromboembolism may be associated with occult cancer.¹⁰ Therefore venous thromboembolism, especially if unprovoked and in older patients, can initiate a malignancy screen, although this practice is not endorsed in current guidelines because of a lack of proven benefit in large randomized trials.^{11,12}

Conversely, the association between arterial thromboembolism and occult cancer is less certain, and while there are numerous case reports describing ischemic stroke or myocardial infarction

serving as the initial presentation of cancer, there are few large-scale, population-based studies detailing the risks of arterial thromboembolism in the months before the pathological diagnosis of cancer.¹³⁻¹⁷ Therefore, we performed a retrospective, matched case-control study evaluating the risks of arterial thromboembolic events in the year before cancer detection. Because there may be a protracted time period of biological activity between the development of cancer and its recognition, we hypothesized that the risks of arterial thromboembolic events increase several months before the pathological diagnosis of cancer.

Materials and Methods

Design

We used linked data from the Surveillance, Epidemiology, and End Results (SEER) registry and Medicare claims from 2004 through 2013 to evaluate the risks of arterial thromboembolic events in the year before a diagnosis of cancer. The SEER-Medicare program comprises multiple population-based cancer registries in the United States, which are linked to Medicare enrollment and claims files, providing detailed clinical information about a diverse population of cancer patients.¹⁸ SEER includes approximately 28% of all patients diagnosed with cancer in the United States.¹⁸ It also provides data from a 5% random sample of Medicare beneficiaries without cancer who live in SEER geographic areas, which enabled us to compare risks of arterial thromboembolic events between patients with incident cancer and control patients without cancer.¹⁸ The Medicare claims data used for this analysis included the physician and supplier file, the outpatient standard analytical file, and the provider analysis and review file. Babak Navi

had full access to all the data in the study and takes responsibility for its integrity and the data analysis. The Memorial Sloan Kettering institutional review board approved this study as exempt research and waived the need for informed consent.

Population

SEER data is organized, and accessed, by individual cancer site making it impractical to analyze all cancer patients registered in SEER. Therefore, for this analysis, we restricted our cancer population to patients \geq 67 years of age diagnosed with breast, lung, prostate, colorectal, bladder, non-Hodgkin lymphoma, uterine, pancreatic, or gastric cancers from January 1, 2005 to December 31, 2013. These cancer sites were selected because they include the 5 most common (overall and solid tumor) cancers, the most common hematological cancer, the most common gynecological cancer, and the 2 cancers historically most associated with thromboembolism (pancreatic and gastric).^{2,19} In total, these cancers account for approximately two-thirds of all incident cancer diagnoses in the United States.¹ If patients were diagnosed first. We excluded patients younger than 67 years of age because most persons in the United States first become eligible for Medicare once they turn 65 and we required sufficient time to evaluate comorbidities in the 730-365 days before study entry.

As done previously, we used site record definitions from the SEER Patient Entitlement and Diagnosis Summary File, as well as the ICD-O-3 site recode classification, to identify the cancer population.^{2,4} The specific site definitions and recodes used were: *breast* (site C500-509; recode

26000); *lung* (sites C300-C301, C310-C319, C320-C329, C339, C340-C349, C381-383, C384, C388, C390, C398, C399; recodes 22010, 22020, 22030, 22050, 22060); *prostate* (site C619; recode 28010); *colorectal* (sites C180-C189, C199, C209, C210-C212, C218, C260; recodes 21041-21049, 21051, 21052, 21060); *bladder* (sites C670-679; recode 29010); *non-Hodgkin lymphoma* (sites C024, C098, C099, C111, C142, C379, C422, C770-779; recodes 33041, 33042); *uterine* (sites C540-549; recodes 27020, 27030); *pancreas* (sites C250-259; recode 21100); and *gastric* (sites C160-C169; recode 21020).

Cancer patients were excluded if they lacked Part A or B Medicare coverage, belonged to a health maintenance organization in the 2 years before their cancer diagnosis, their month of cancer diagnosis was missing, their age at the time of cancer diagnosis was <67 years, or we could not identify a control patient without cancer to match. In order to minimize misclassification error and to focus on first arterial thromboembolic events, we also excluded patients if they had an International Classification of Diseases-9-Clinical Modification (ICD-9-CM) diagnosis code for cerebrovascular disease (430.xx-438.xx) or coronary heart disease (410.xx-414.xx) 730-365 days before their date of cancer diagnosis.

All cancer patients were individually matched to a control patient without cancer by year of birth, sex, race (white or nonwhite), SEER registry (surrogate for geographic region), Charlson comorbidity index (trichotomized into 0, 1, and ≥ 2), and the diagnoses of hypertension (ICD-9-CM codes 401.xx-405.xx, 437.2) and atrial fibrillation (ICD-9-CM codes 427.31, 427.32) in the 730-365 days before study entry; these latter diagnoses were individually matched upon because they are important cardiovascular risk factors missing from the Charlson comoribidity index.²⁰

Similar to cancer cases, we excluded control patients if they lacked Medicare Part A or B coverage, belonged to a health maintenance organization in the 2 years before study entry, or they had a Medicare claim for cerebrovascular or coronary heart disease 730-365 days before their date of study entry.

Measurements

We defined an arterial thromboembolic event as the composite of myocardial infarction or ischemic stroke. We also assessed myocardial infarction alone and ischemic stroke alone. Validated ICD-9-CM diagnostic code algorithms were used to identify these events.^{21,22} Myocardial infarction was identified by inpatient code 410.xx in any diagnostic position.²¹ Ischemic stroke was identified by inpatient codes 433.x1, 434.x1, or 436 in any diagnostic position without concurrent codes for rehabilitation (V57) in the primary diagnostic position, or trauma (800.xx-804.xx, 850.xx-854.xx) or hemorrhagic stroke (430.xx, 431.xx) in any diagnostic position.²² These diagnostic code algorithms have been previously shown to have \geq 90% positive predictive value for correctly identifying myocardial infarction and ischemic stroke when compared to detailed chart review.^{21,22} Mesenteric ischemia and systemic artery thromboembolism were not ascertained for the primary analysis because these are rare diseases that lack reliable diagnostic codes for identification through claims data.²³

Statistical Analysis

Descriptive statistics with exact confidence intervals were used to describe patient characteristics at the time of study entry, which was defined as the date of cancer diagnosis for both cancer patients and their matched controls without cancer. Risks of arterial thromboembolic events were compared between groups during 1-month periods in the 12 months before study entry. The Mantel-Haenszel estimator for matched data was used to calculate odds ratios and absolute risks for each 1-month period.

Several sensitivity analyses were performed. First, to account for the unknown exact date of cancer diagnosis within SEER (only month and year are provided so per standard practice we used the middle of the month for analyses), we excluded events that occurred during the 15 days immediately before the recorded date of cancer diagnosis. Second, to account for the potential confounding effects of withholding anticoagulation for biopsy, we excluded patients with a diagnosis of atrial fibrillation. Third, to account for an imbalance in the rate of chronic obstructive pulmonary disease between groups, we matched cancer patients to cancer-free controls by this diagnosis (ICD-9-CM codes 490.xx-492.xx, 494.xx, 496.xx), in addition to the variables previously matched by in our primary analysis.

In secondary analysis, odds ratios and absolute risks were stratified by cancer stage at diagnosis. As in prior SEER-Medicare studies, cancer staging was defined according to the American Joint Committee on Cancer classification, except for prostate cancer, which was staged according to the T (clinical) classification, and non-Hodgkin lymphoma, which was staged according to the Ann Arbor classification.^{2,24-26} Patients with unknown cancer stage (and their matched controls) were excluded from this secondary analysis.

In a post-hoc exploratory analysis, we expanded the arterial thromboembolic event composite outcome to include systemic artery thromboembolism and acute mesenteric ischemia, in addition to acute myocardial infarction and ischemic stroke. Systemic artery thromboembolism was identified by inpatient code 444.xx in any diagnostic position. Acute mesenteric ischemia was identified by inpatient code 557.0 in any diagnostic position. An alpha error of 5% was used to determine statistical significance. All analyses were performed by A.S.R. using SAS (version 9.4, SAS Institute Inc.) or R (version 3.4.3) software.

Data Sharing Statement

The data used in this analysis includes restricted SEER-Medicare claims, which based on the terms of the data use agreement, cannot be shared directly with other investigators. However, investigators can obtain access to these data by application to the National Cancer Institute.

Results

Baseline Characteristics

We identified 374,331 pairs of cancer patients and matched controls without cancer who met study eligibility criteria. The mean age was 76 years (standard deviation 7) and 52% were women (Table 1). Prostate (n=82,099), breast (n=74,638), lung (n=74,633), and colorectal

(n=56,366) cancers accounted for 77% of cancer types. At diagnosis, 30% of cancers were stages 3 or 4.

Primary Analysis

From 360 to 151 days before cancer diagnosis, the 30-day interval risks of arterial thromboembolic events were similar between cancer patients and matched controls without cancer (Figure 1, Table 2). From 150 to 1 day before cancer diagnosis, the interval 30-day risks of arterial thromboembolic events were higher in cancer patients versus matched controls, progressively increasing as the cancer diagnosis date approached, and peaking during the 30days immediately before cancer diagnosis, when 2,313 (0.62%) cancer patients were diagnosed with an arterial thromboembolic event versus 413 (0.11%) cancer-free controls (odds ratio [OR], 5.63; 95% confidence interval [CI], 5.07-6.25; p<0.001). When analyzing all 360 days before study entry, 6,567 (1.75%) cancer patients were diagnosed with an arterial thromboembolic event versus 3,916 (1.05%) cancer-free controls (OR, 1.69; 95% CI, 1.63-1.76; p<0.001). Among cancer patients who had an arterial thromboembolic event in the 360 days before cancer 24%), prostate (n=752, 11%), breast (n=663, 10%), bladder (n=533, 8%), non-Hodgkin lymphoma (n=384, 6%), pancreatic (n=333, 5%), gastric (n=250, 4%), and uterine (n=183, 3%) (Table 3).

Sensitivity Analysis

When excluding events that occurred during the 15 days immediately before the recorded date of cancer diagnosis (i.e., analysis restricted to 16-360 days before study entry), risk differences between groups were attenuated but still significant with 4,814 (1.29%) arterial thromboembolic events diagnosed in cancer patients versus 3,718 (0.99%) events in cancer-free controls (OR, 1.31; 95% CI, 1.25-1.37; p<0.001). In the 16-30 days before study entry, 561 (0.15%) cancer patients were diagnosed with an arterial thromboembolic event versus 215 (0.06%) cancer-free controls (OR, 2.62; 95% CI, 2.24-3.07; p<0.001).

When restricted to patients without a diagnosis of atrial fibrillation (357,905 pairs of patients), in the 360 days before study entry, 6,190 (1.73%) cancer patients were diagnosed with an arterial thromboembolic event versus 3,633 (1.02%) cancer-free controls. In these patients, the risk of an arterial thromboembolic event first became increased 150 days before the date of cancer diagnosis, and thereafter progressively increased until peaking in the 30 days preceding the diagnosis (OR, 5.77; 95% CI, 5.18-6.44).

When cancer patients and their cancer-free controls were matched by the diagnosis of chronic obstructive pulmonary disease in addition to the variables previously matched by in our primary analysis (298,459 pairs of patients), there remained an increased risk of arterial thromboembolic events in the 360 days before the date of cancer diagnosis (OR, 1.91; 95% CI, 1.82-2.00; p<0.001). This risk first became consistently increased at 150 days before the date of cancer diagnosis, and thereafter progressively increased until peaking in the 30 days preceding the diagnosis (OR, 6.30; 95% CI, 5.56-7.14).

Secondary Analyses

Cancer patients' relative risks for myocardial infarction and ischemic stroke in isolation were similar, although their absolute risk for myocardial infarction appeared slightly higher than for ischemic stroke (Table 4). The risks of both myocardial infarction and ischemic stroke were mostly equal between cancer patients and cancer-free controls until about 150 days before cancer diagnosis, when risks became higher among cancer patients and progressively increased until the date of cancer diagnosis. In the 360 days before study entry, there were 3,877 (1.04%)myocardial infarctions diagnosed in cancer patients versus 2,171 (0.58%) in cancer-free controls (OR, 1.80; 95% CI, 1.71-1.90; p<0.001). Meanwhile, during this 360 day period, there were 2,913 (0.78%) ischemic strokes diagnosed in cancer patients versus 1,836 (0.49%) in cancer-free controls (OR, 1.59; 95% CI, 1.50-1.69; p<0.001).

Stage Analyses

Among cancer patients who had an arterial thromboembolic event within 360 days of cancer diagnosis, the cancer stage at diagnosis was stage 0/1 in 1,871 (28%), stage 2 in 1,065 (16%), stage 3 in 1,044 (16%), stage 4 in 1,596 (24%), and an unknown stage in 991 (15%). Higher cancer stages at cancer diagnosis were associated with increased odds of an arterial thromboembolic event. For instance, compared to cancer-free controls, cancer patients' relative odds for an arterial thromboembolic event in the 360 days before cancer diagnosis was 1.32 (95% CI, 1.22-1.43) for stage 1 cancers, 1.55 (95% CI, 1.40-1.70) for stage 2 cancers, 2.19 (95%

CI, 1.96-2.44) for stage 3 cancers, and 2.17 (95% CI, 1.99-2.37) for stage 4 cancers. These analyses were similar when evaluating myocardial infarction alone and ischemic stroke alone.

Exploratory Analysis

When using an expanded definition of arterial thromboembolic events that included systemic artery thromboembolism and acute mesenteric ischemia, in the 360 days before study entry, 7,263 (1.94%) cancer patients were diagnosed with an arterial thromboembolic event versus 4,215 (1.13%) cancer-free control patients (OR, 1.74; 95% CI, 1.68-1.81; p<0.001). This included 491 (0.13%) systemic artery thromboembolism and 304 (0.08%) acute mesenteric ischemia diagnoses among cancer patients and 209 (0.06%) systemic artery thromboembolism and 127 (0.03%) acute mesenteric ischemia diagnoses among cancer patients diagnoses among cancer-free controls. The risk of the expanded arterial thromboembolic event outcome was highest in the 30 days immediately before the cancer diagnosis (OR, 6.03; 95% CI, 5.45-6.67).

Discussion

In an analysis of over 700,000 Medicare beneficiaries, we found that in the 360 days before a cancer diagnosis, the risk of an arterial thromboembolic event was increased nearly 70%. This risk first became increased about 5 months before cancer diagnosis and thereafter progressively rose until peaking in the month before cancer diagnosis, when the risk was increased more than fivefold. The relative risks of myocardial infarction alone and ischemic stroke alone were similar, although myocardial infarction was slightly more common than ischemic stroke. Of the

From www.bloodjournal.org by guest on December 21, 2018. For personal use only.

cancer types studied, lung and colorectal cancer were the most likely cancers to be preceded by an arterial thromboembolic event. Meanwhile, among cancer patients who had preceding arterial thromboembolic events, 40% had stages 3 or 4 disease upon cancer diagnosis.

Few population-based studies have systematically analyzed the association between cancer and preceding arterial thromboembolic events. Among 3,680 adults with nondisabling ischemic stroke who participated in a randomized trial of vitamin supplements, the 1-year cancer incidence rate was 1.2%, and this was increased 20% compared to the estimated age-adjusted rate in the general population.¹⁵ In a prospective Norwegian study of 28,763 patients, incident myocardial infarction was associated with a 46% increased incidence of cancer during followup.¹⁶ Additionally, in the first 6 months after myocardial infarction, the risk of cancer was increased 120%.¹⁶ In a retrospective analysis of 6,600 Danish patients with lower limb arterial thrombosis, the incidence rate of subsequent cancer diagnoses was increased 35% compared to expected national rates, and the highest relative risk was in the first 6-months after arterial thrombosis, when it was more than tripled.¹⁷ Our study expands on these analyses by including a more heterogeneous patient population, a matched study design whereby cancer patients were matched to cancer-free controls by demographics and comorbidities to lower the risk of confounding, and a substantially larger sample size with resultant increased statistical power, which enabled us to delineate risks by monthly intervals.

Our findings also build on previous studies that evaluated the risk of arterial thromboembolic events after cancer diagnosis. In our prior SEER-Medicare study of 279,719 patients with 8 common cancers, the risk for an arterial thromboembolic event in the 6 months after cancer

diagnosis was increased 120% compared to matched controls without cancer.² Additionally, similar to the current study, risks were higher closer to the time of cancer diagnosis, peaking in the first month afterwards, when the hazard ratio for an arterial thromboembolic event was 5.2.² Meanwhile, nationwide Swedish studies that evaluated over 820,000 cancer patients, including cancer types not included in the aforementioned SEER-Medicare study, reported analagous findings, with most cancers conferring a transient increased risk of stroke and coronary heart disease that attenuated or disappeared within 1 year of cancer diagnosis.^{5,6} By demonstrating that arterial thromboembolism risk increases not only after diagnosis but also several months before, our study suggests that much of the increased risk may be due to the cancer's intrinsic effects on coagulation, rather than extrinsic factors that only develop after cancer diagnosis, such as cancer treatment effects and antithrombotic interruption for chemotherapy-induced thrombocytopenia. However, antithrombotic interruption for procedures, especially surgical biopsy of a newly discovered mass, might have contributed to the increased arterial thromboembolism risk before cancer diagnosis, particularly in the month or two before.

This study has several important limitations. First, it was retrospective and relied on Medicare claims data for arterial thromboembolic event diagnoses and therefore misclassification error is possible. We attempted to minimize this concern by using previously validated diagnostic code algorithms to identify events; however, it remains possible that some diagnoses were incorrect, and that other conditions, including cancer metastases, mimicked thromboembolic events. Similarly, by using claims data, we lacked clinical information on lifestyle factors such as smoking, examination and laboratory data, pathophysiological mechanisms, severity of events, and electrocardiographic and imaging findings. We also lacked reliable data on the use of

antithrombotic therapy; hence, we were unable to determine whether its interruption for surgical biopsy contributed to arterial thromboembolism risk. Second, because this was a SEER-Medicare study, we restricted our cohort to patients 67 years of age and older, so our results may not apply to younger patients; although in the United States, the estimated median age of patients with cancer is 66 and the estimated mean ages of patients with myocardial infarction and ischemic stroke is 69 and 69, respectively.²⁷⁻²⁹ Third, since cancer patients were matched to cancer-free controls, only those who survived an arterial thromboembolic event and lived long enough to be diagnosed with cancer were included, and therefore immortal time bias could have affected the association between cancer and preceding arterial thromboembolic events if cancer patients and controls differed in their likelihood to survive events. However, the majority of myocardial infarctions and ischemic strokes are nonfatal and, in practical terms, an association between two diseases is most relevant when patients survive the first disease to become at risk for the second.³⁰ Fourth, some arterial thromboembolic events may have led to a diagnosis of cancer through increased medical surveillance. For instance, a chest x-ray performed during myocardial infarction hospitalization might demonstrate a lung mass, which is later biopsied and determined to be cancer. This phenomenon would be most plausible in the month preceding cancer diagnosis and may partly explain the very high odds of thromboembolic events during that month, although some myocardial infarctions probably led to drug-eluting stent placement, which could have delayed cancer diagnosis by several months because of the need for dual antiplatelets to prevent in-stent thrombosis. Fifth, because SEER organizes and releases data by cancer site, we were unable to evaluate all cancer patients registered in SEER. However, we did include 9 common cancer sites, including the 5 most common solid tumor cancers and the most common hematological cancer, which combined, account for approximately two-thirds of all

cancer in the United States.¹ Finally, while we matched on multiple variables, and there were similar rates of unmatched cardiovascular and cancer risk factors between groups, it is possible that residual confounding biased our results. Smoking history, in particular, is not reliably identified through claims data, and any potential imbalances in smoking rates could have confounded the association between cancer and preceding arterial thromboembolic events. However, if present, confounding bias would be expected to produce uniform risk differences between groups over time; while in our analysis, risks between groups varied greatly by time.

In summary, the risk of myocardial infarction and ischemic stroke is increased approximately 70% in the year before cancer diagnosis. This increased risk begins 5 months before cancer is diagnosed pathologically and peaks in the month before when it is increased fivefold. Risk of arterial thromboembolism preceding cancer diagnosis varies by primary cancer type and seems highest with lung and colorectal cancers. These data suggest that some myocardial infarction and ischemic stroke events may be triggered, or potentially caused by, occult cancer. Future research is needed to identify clinically useful biomarkers for occult cancer in patients with arterial thromboembolic events and to determine the utility of cancer screening strategies in these patients, particularly among those with cancer risk factors or unexplained "cryptogenic" events. In the meantime, we recommend that patients with acute myocardial infarction and ischemic stroke be up-to-date with their age- and gender-appropriate cancer screening, and that their clinicians pay close attention to, and have a low threshold to investigate, any symptoms or signs consistent with occult cancer, such as unexplained anemia or weight loss.

Acknowledgments

This study was supported by research funding from the NIH through grants K23NS091395 (B.B.N.) and P30CA008748 (A.S.R., K.S.P., and L.M.D.), and the Florence Gould Endowment for Discovery in Stroke (B.B.N.). The sponsors had no role in the design and/or execution of this study. The authors thank Monica Chen for her copyediting and clerical support and for assisting in the design and creation of the visual abstract.

Authorship

Contribution: B.B.N., A.S.R., H.K., K.S.P., and L.M.D. conceived the study and its design; B.B.N. and K.S.P. acquired the data; B.B.N., A.S.R., H.K., C.I., P.M.O., S.T.T., K.S.P., and L.M.D. analyzed and interpreted the data; B.B.N. drafted the manuscript; B.B.N., A.S.R., H.K., C.I., P.M.O., S.T.T., K.S.P., and L.M.D. revised the manuscript for important intellectual content; B.B.N., A.S.R., and K.S.P. performed the statistical analysis.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

Correspondence: Babak B. Navi, Weill Cornell Medicine, Department of Neurology, 525 East 68th Street, Room F610, New York, NY 10065; ban9003@med.cornell.edu.

References

 Howlader N, Noone AM, Krapcho M, et al. SEER Cancer Statistics Review, 1975-2014, National Cancer Institute. Available at: <u>http://seer.cancer.gov/csr/1975_2014/</u>. Accessed March 1, 2018.

2. Navi BB, Reiner AS, Kamel H, et al. Risk of Arterial Thromboembolism in Patients With Cancer. *J Am Coll Cardiol*. 2017;70(8):926-938.

3. Yeh ETH, Chang HM. Cancer and Clot: Between a Rock and a Hard Place. *J Am Coll Cardiol.* 2017;70(8):939-941.

4. Navi BB, Reiner AS, Kamel H, et al. Association between incident cancer and subsequent stroke. *Ann Neurol*. 2015;77(2):291-300.

5. Zoller B, Ji J, Sundquist J, Sundquist K. Risk of coronary heart disease in patients with cancer: a nationwide follow-up study from Sweden. *Eur J Cancer*. 2012;48(1):121-128.

6. Zoller B, Ji J, Sundquist J, Sundquist K. Risk of haemorrhagic and ischaemic stroke in patients with cancer: a nationwide follow-up study from Sweden. *Eur J Cancer*.

2012;48(12):1875-1883.

7. Bang OY, Chung JW, Lee MJ, et al. Cancer Cell-Derived Extracellular Vesicles Are Associated with Coagulopathy Causing Ischemic Stroke via Tissue Factor-Independent Way: The OASIS-CANCER Study. *PLoS One*. 2016;11(7):e0159170.

8. Demers M, Wagner DD. NETosis: a new factor in tumor progression and cancerassociated thrombosis. *Semin Thromb Hemost*. 2014;40(3):277-283.

9. Falanga A, Marchetti M, Vignoli A. Coagulation and cancer: biological and clinical aspects. *J Thromb Haemost*. 2013;11(2):223-233.

10. Carrier M, Le Gal G, Wells PS, Fergusson D, Ramsay T, Rodger MA. Systematic review: the Trousseau syndrome revisited: should we screen extensively for cancer in patients with venous thromboembolism? *Ann Intern Med.* 2008;149(5):323-333.

11. Khorana AA, Carrier M, Garcia DA, Lee AY. Guidance for the prevention and treatment of cancer-associated venous thromboembolism. *J Thromb Thrombolysis*. 2016;41(1):81-91.

12. Carrier M, Lazo-Langner A, Shivakumar S, et al. Screening for Occult Cancer in Unprovoked Venous Thromboembolism. *N Engl J Med*. 2015;373(8):697-704.

13. Taccone FS, Jeangette SM, Blecic SA. First-ever stroke as initial presentation of systemic cancer. *J Stroke Cerebrovasc Dis*. 2008;17(4):169-174.

14. Navi BB, DeAngelis LM, Segal AZ. Multifocal strokes as the presentation of occult lung cancer. *J Neurooncol*. 2007;85(3):307-309.

15. Qureshi AI, Malik AA, Saeed O, Adil MM, Rodriguez GJ, Suri MF. Incident cancer in a cohort of 3,247 cancer diagnosis free ischemic stroke patients. *Cerebrovasc Dis*. 2015;39(5-6):262-268.

16. Rinde LB, Smabrekke B, Hald EM, et al. Myocardial infarction and future risk of cancer in the general population-the Tromso Study. *Eur J Epidemiol*. 2017;32(3):193-201.

17. Sundboll J, Veres K, Horvath-Puho E, Adelborg K, Sorensen HT. Risk and Prognosis of Cancer After Lower Limb Arterial Thrombosis. *Circulation*. 2018.

Howlader N, Noone AM, Krapcho M, et al. Surveillance Epidemiology and End Results
 (SEER) Program. SEER-Medicare Linked Database. National Cancer Institute.

http://healthcaredelivery.cancer.gov/seermedicare/. Accessed March 21, 2018.

Khorana AA, Kuderer NM, Culakova E, Lyman GH, Francis CW. Development and validation of a predictive model for chemotherapy-associated thrombosis. *Blood*.
 2008;111(10):4902-4907.

20. Charlson ME, Pompei P, Ales KL, MacKenzie CR. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis*. 1987;40(5):373-383.

21. Kiyota Y, Schneeweiss S, Glynn RJ, Cannuscio CC, Avorn J, Solomon DH. Accuracy of Medicare claims-based diagnosis of acute myocardial infarction: estimating positive predictive value on the basis of review of hospital records. *Am Heart J*. 2004;148(1):99-104.

22. Tirschwell DL, Longstreth WT, Jr. Validating administrative data in stroke research. *Stroke*. 2002;33(10):2465-2470.

23. Korabathina R, Weintraub AR, Price LL, et al. Twenty-year analysis of trends in the incidence and in-hospital mortality for lower-extremity arterial thromboembolism. *Circulation*. 2013;128(2):115-121.

24. Edge SB, Compton CC. The American Joint Committee on Cancer: the 7th edition of the AJCC cancer staging manual and the future of TNM. *Ann Surg Oncol.* 2010;17(6):1471-1474.

Carbone PP, Kaplan HS, Musshoff K, Smithers DW, Tubiana M. Report of the
 Committee on Hodgkin's Disease Staging Classification. *Cancer Res.* 1971;31(11):1860-1861.

26. Chamie K, Williams SB, Hershman DL, Wright JD, Nguyen PL, Hu JC. Populationbased assessment of determining predictors for quality of prostate cancer surveillance. *Cancer*. 2015;121(23):4150-4157.

27. Kissela BM, Khoury JC, Alwell K, et al. Age at stroke: temporal trends in stroke incidence in a large, biracial population. *Neurology*. 2012;79(17):1781-1787.

28. Yeh RW, Sidney S, Chandra M, Sorel M, Selby JV, Go AS. Population trends in the incidence and outcomes of acute myocardial infarction. *N Engl J Med*. 2010;362(23):2155-2165.

29. NIH National Cancer Institute. Age and Cancer Risk. <u>https://www.cancer.gov/about-</u> cancer/causes-prevention/risk/age. Accessed September 3.

30. Benjamin EJ, Virani SS, Callaway CW, et al. Heart Disease and Stroke Statistics-2018

Update: A Report From the American Heart Association. Circulation. 2018;137(12):e67-e492.

Table 1. Characteristics of Pairs of Cancer Patients and Matched Controls, 2004-2013					
	Cancer Patients	Controls without Cancer			
Characteristic	$(N=374,331)^*$	(N=374,331)			
Age, mean (SD), y	76 (7)	76 (7)			
Sex					
Women	193,726 (52)	193,726 (52)			
Men	180,605 (48)	180,605 (48)			
Race					
White	314,387 (84)	314,387 (84)			
Non-white	59,944 (16)	59,944 (16)			
Geographic region					
Northeast	75,013 (20)	75,013 (20)			
Midwest	61,900 (17)	61,900 (17)			
South	96,606 (26)	96,606 (26)			
West	140,812 (38)	140,812 (38)			
Charlson Comorbidities					
0	260,814 (70)	260,814 (70)			
1	79,136 (21)	79,136 (21)			
2 or more	34,381 (9)	34,381 (9)			
Hypertension	221,127 (59)	221,127 (59)			
Atrial fibrillation	16,426 (4)	16,426 (4)			
Diabetes mellitus	92,915 (25)	96,338 (26)			
Peripheral vascular disease	36,444 (10)	35,463 (9)			
COPD	54,300 (15)	42,736 (11)			
Congestive heart failure	15,458 (4)	15,537 (4)			
Cardiac valvular disease	27,287 (7)	29,352 (8)			
Liver disease	1,481 (0.4)	1,592 (0.4)			
Chronic kidney disease	14,218 (4)	14,405 (4)			
Census tract poverty level <10% [†]	182,239 (49)	NA			
Cancer type	· · · · · ·				
Breast	74,638 (20)	NA			
Lung	74,633 (20)	NA			
Prostate	82,099 (22)	NA			
Colorectal	56,366 (15)	NA			
Bladder	26,503 (7)	NA			
Non-Hodgkin lymphoma	21,560 (6)	NA			
Uterine	13,473 (4)	NA			
Panreatic	16,386 (4)	NA			
Gastric	8,673 (2)	NA			
Cancer stage [‡]					
0	26,685 (7)	NA			
1	115,783 (31)	NA			
2	72,636 (19)	NA			
3	44,854 (12)	NA			
4	67,121 (18)	NA			
Unknown	47,252 (13)	NA			

Abbreviations: SD, standard deviation; COPD, chronic obstructive pulmonary disease; NA, not applicable.

*Data presented as number (%) unless otherwise specified. Due to rounding, some values don't add to 100.

[†]Number and proportion of patients who live in areas where <10% of people are below the poverty line according to the

From www.bloodjournal.org by guest on December 21, 2018. For personal use only.

2000 census.

[‡]Refers to the American Joint Committee on Cancer (AJCC) sixth edition staging schema, except for patients with prostate cancer, who were staged according to the T (clinical) staging classification and patients with non-Hodgkin lymphoma, who were staged according to the Ann Arbor staging classification.

Table 2. Risk of Arterial Thromboembolic Events [*] Prior to Cancer Diagnosis						
	Cancer Patients	Cancer-free Controls				
Time Period Before	No. and Percentage	No. and Percentage	Odds Ratio and	P-value		
Cancer Diagnosis	of Events	of Events	95% CI			
1-30 days	2,313 (0.62%)	413 (0.11%)	5.63 (5.07-6.25)	< 0.001		
31-60 days	635 (0.17%)	365 (0.10%)	1.74 (1.53-1.98)	< 0.001		
61-90 days	496 (0.13%)	315 (0.08%)	1.58 (1.37-1.81)	< 0.001		
91-120 days	431 (0.12%)	349 (0.09%)	1.24 (1.07-1.42)	0.003		
121-150 days	427 (0.11%)	342 (0.09%)	1.25 (1.08-1.44)	0.002		
151-180 days	379 (0.10%)	333 (0.09%)	1.14 (0.98-1.32)	0.08		
181-210 days	312 (0.08%)	291 (0.08%)	1.07 (0.91-1.26)	0.39		
211-240 days	370 (0.10%)	318 (0.08%)	1.17 (1.00-1.35)	0.05		
241-270 days	309 (0.08%)	306 (0.08%)	1.01 (0.86-1.18)	0.90		
271-300 days	316 (0.08%)	311 (0.08%)	1.02 (0.87-1.19)	0.84		
301-330 days	334 (0.09%)	313 (0.08%)	1.07 (0.92-1.25)	0.41		
331-360 days	317 (0.08%)	300 (0.08%)	1.06 (0.90-1.24)	0.49		
Abbreviations: No., number: CI, confidence interval.						

*Composite of myocardial infarction and ischemic stroke diagnoses.

Table 3. Characteristics of Cancer Patients Stratified by the Diagnosis of Arterial Thromboembolic Events in the Year Before Cancer Diagnosis

Characteristic	Preceding Arterial Thromboembolic Event (N=6 567) [*]	No Preceding Arterial Thromboembolic Event (N=367 764)
Age, mean (SD), v	79 (7)	76 (7)
Sex Sex		
Women	3,308 (50)	190,418 (52)
Men	3,259 (50)	177,346 (48)
Race	- , (/	
White	5,402 (82)	308,985 (84)
Non-white	1,165 (18)	58,779 (16)
Geographic region		· · · · · · · · · · · · · · · · · · ·
Northeast	1,522 (23)	73,491 (20)
Midwest	1,061 (16)	60,839 (17)
South	1,844 (28)	94,762 (26)
West	2,140 (33)	138,672 (38)
Charlson Comorbidities		
0	4,121 (63)	256,693 (70)
1	1,480 (23)	77,656 (21)
2 or more	966 (15)	33,415 (9)
Hypertension	3,836 (58)	217,291 (59)
Atrial fibrillation	377 (6)	16,049 (4)
Diabetes mellitus	1,795 (27)	91,120 (25)
Peripheral vascular disease	917 (14)	35,527 (10)
COPD	1,183 (18)	53,117 (14)
Congestive heart failure	482 (7)	14,976 (4)
Cardiac valvular disease	533 (8)	26,754 (7)
Liver disease	20 (0.3)	1,461 (0.4)
Chronic kidney disease	386 (6)	13,832 (4)
Census tract poverty level <10% [†]	2,895 (44)	179,344 (49)
Cancer type		
Breast	663 (10)	73,975 (20)
Lung	1,908 (29)	72,725 (20)
Prostate	752 (11)	81,347 (22)
Colorectal	1,561 (24)	54,805 (15)
Bladder	533 (8)	25,970 (7)
Non-Hodgkin lymphoma	384 (6)	21,176 (6)
Uterine	183 (3)	13,290 (4)
Pancreatic	333 (5)	16,053 (4)
Gastric	250 (4)	8,423 (2)
Cancer stage [‡]		
0	364 (6)	26,321 (7)
1	1,507 (23)	114,276 (31)
2	1,065 (16)	71,571 (19)
3	1,044 (16)	43,810 (12)
4	1,596 (24)	65,525 (18)
Unknown	991 (15)	46,261 (13)

Abbreviations: SD, standard deviation; COPD, chronic obstructive pulmonary disease; NA, not applicable.

*Data presented as number (%) unless otherwise specified. Due to rounding, some values don't add to 100.

[†]Number and proportion of patients who live in areas where <10% of people are below the poverty line according to the 2000 census.

[‡]Refers to the American Joint Committee on Cancer (AJCC) sixth edition staging schema, except for patients with prostate cancer, who were staged according to the T (clinical) staging classification and patients with non-Hodgkin lymphoma, who were staged according to the Ann Arbor staging classification.

Table 4. Risks of Myocardial Infarction and Ischemic Stroke Prior to Cancer Diagnosis						
	Cancer Patients	Cancer-free Controls				
Time Period Before	No. and Percentage	No. and Percentage	Odds Ratio and	P-value		
Cancer Diagnosis	of Outcomes	of Outcomes	95% CI			
Myocardial Infarctio	n					
1-30 days	1,408 (0.38%)	220 (0.06%)	6.43 (5.57-7.41)	< 0.001		
31-60 days	340 (0.09%)	201 (0.05%)	1.69 (1.42-2.01)	< 0.001		
61-90 days	274 (0.07%)	170 (0.05%)	1.61 (1.33-1.95)	< 0.001		
91-120 days	241 (0.06%)	187 (0.05%)	1.29 (1.07-1.56)	0.009		
121-150 days	237 (0.06%)	192 (0.05%)	1.24 (1.02-1.50)	0.03		
151-180 days	222 (0.06%)	200 (0.05%)	1.11 (0.92-1.34)	0.28		
181-210 days	170 (0.05%)	170 (0.05%)	1.00 (0.81-1.24)	1.00		
211-240 days	222 (0.06%)	167 (0.04%)	1.33 (1.09-1.63)	0.005		
241-270 days	185 (0.05%)	162 (0.04%)	1.14 (0.93-1.41)	0.22		
271-300 days	199 (0.05%)	171 (0.05%)	1.16 (0.95-1.43)	0.15		
301-330 days	196 (0.05%)	155 (0.04%)	1.27 (1.02-1.56)	0.03		
331-360 days	183 (0.05%)	176 (0.05%)	1.04 (0.85-1.28)	0.71		
Ischemic Stroke						
1-30 days	979 (0.26%)	195 (0.05%)	5.04 (4.32-5.88)	< 0.001		
31-60 days	307 (0.08%)	170 (0.05%)	1.81 (1.50-2.18)	< 0.001		
61-90 days	232 (0.06%)	154 (0.04%)	1.51 (1.23-1.85)	< 0.001		
91-120 days	195 (0.05%)	169 (0.05%)	1.15 (0.94-1.42)	0.17		
121-150 days	196 (0.05%)	153 (0.04%)	1.28 (1.04-1.58)	0.02		
151-180 days	168 (0.04%)	141 (0.04%)	1.19 (0.95-1.49)	0.13		
181-210 days	144 (0.04%)	122 (0.03%)	1.18 (0.93-1.50)	0.18		
211-240 days	156 (0.04%)	157 (0.04%)	0.99 (0.80-1.24)	0.95		
241-270 days	126 (0.03%)	144 (0.04%)	0.88 (0.69-1.11)	0.27		
271-300 days	128 (0.03%)	144 (0.04%)	0.89 (0.70-1.13)	0.33		
301-330 days	144 (0.04%)	162 (0.04%)	0.89 (0.71-1.11)	0.30		
331-360 days	138 (0.04%)	125 (0.03%)	1.10 (0.87-1.41)	0.42		
Abbreviations: No., number; CI, confidence interval.						

Figure Legend

Figure 1. Relative and absolute odds of an arterial thromboembolic event during 30-day intervals in the 360 days before cancer diagnosis. The comparator group for the relative odds calculation were control patients without cancer who were individually matched to cancer cases by age, sex, race, geographic region, Charlson comorbidity index, history of hypertension, and history of atrial fibrillation. Arterial thromboembolic events were defined as a composite of myocardial infarction and ischemic stroke. Dashed blue lines indicate 95% confidence intervals for the relative odds of arterial thromboembolic events.



Absolute 30-Day Rates of Arterial Thromboembolism (%)



Prepublished online December 21, 2018; doi:10.1182/blood-2018-06-860874

Arterial thromboembolic events preceding the diagnosis of cancer in older persons

Babak B. Navi, Anne S. Reiner, Hooman Kamel, Costantino Iadecola, Peter M. Okin, Scott T. Tagawa, Katherine S. Panageas and Lisa M. DeAngelis

Information about reproducing this article in parts or in its entirety may be found online at: http://www.bloodjournal.org/site/misc/rights.xhtml#repub_requests

Information about ordering reprints may be found online at: http://www.bloodjournal.org/site/misc/rights.xhtml#reprints

Information about subscriptions and ASH membership may be found online at: http://www.bloodjournal.org/site/subscriptions/index.xhtml

Advance online articles have been peer reviewed and accepted for publication but have not yet appeared in the paper journal (edited, typeset versions may be posted when available prior to final publication). Advance online articles are citable and establish publication priority; they are indexed by PubMed from initial publication. Citations to Advance online articles must include digital object identifier (DOIs) and date of initial publication.

Blood (print ISSN 0006-4971, online ISSN 1528-0020), is published weekly by the American Society of Hematology, 2021 L St, NW, Suite 900, Washington DC 20036. Copyright 2011 by The American Society of Hematology; all rights reserved.